arXiv:1510.05036v1 [math.FA] 16 Oct 2015

MISCELLANOUS APPLICATIONS OF CERTAIN MINIMAX THEOREMS. I

B. RICCERI

Department of Mathematics, University of Catania
Viale A. Doria 6, 95125 Catania, Italy

ABSTRACT. Here is one of the results of this paper (with the convention $\frac{1}{0}=+\infty$): Let X be a real Hilbert space and let $J:X\to\mathbf{R}$ be a C^1 functional, with compact derivative, such that

$$\alpha^*:=\max\left\{0,\ \limsup_{\|x\|\to+\infty}\frac{J(x)}{\|x\|^2}\right\}<\beta^*:=\sup_{x\in X\setminus\{0\}}\frac{J(x)}{\|x\|^2}<+\infty\ .$$

Then, for every $\lambda\in\left]\frac{1}{2\beta^*},\frac{1}{2\alpha^*}\right[$ and for every convex set $C\subseteq X$ dense in X, there exists $\tilde{y}\in C$ such that the equation

$$x=\lambda J'(x)+\tilde{y}$$

has at least three solutions, two of which are global minima of the functional $x\to\frac{1}{2}\|x\|^2-\lambda J(x)-\langle x,\tilde{y}\rangle$.

AMS (MOS) Subject Classification. 49K35; 90C47; 47G40; 35A15; 35J20

1. RESULTS

The present one is the first of a series of notes (with the same title) on new consequences and applications of certain minimax theorems that we have established in the past years ([2]-[10]).

In [8], in particular, we obtained the following result:

Theorem A. ([8], Theorem 3.4) *Let X be a non-empty set, Y a real inner product space and $I:X\to\mathbf{R}$, $\Phi:X\to Y$ two given functions.*

Then, for each $\mu>0$, at least one of the following assertions holds:

(a) for each filtering cover $\mathcal{N}$ of X, there exists $A\in\mathcal{N}$ such that

$$\sup_{y\in Y}\inf_{x\in A}(I(x)+\mu(2\langle\Phi(x),y\rangle-\|y\|^2))<\inf_{x\in A}\sup_{y\in\Phi(A)}(I(x)+\mu(2\langle\Phi(x),y\rangle-\|y\|^2))\ ;$$

(b) for each global minimum u of $x\to I(x)+\mu\|\Phi(x)\|^2$, one has

$$I(u)\le I(x)+2\mu(\langle\Phi(x),\Phi(u)\rangle-\|\Phi(u)\|^2)$$

for all $x\in X$.

A cover $\mathcal{N}$ of the set X is said to be filtering if for each $A_1,A_2\in\mathcal{N}$, there exists $A_3\in\mathcal{N}$ such that $A_1\cup A_2\subseteq A_3$.

In the present short note, we want to highlight two applications of the following consequences of Theorem A:

Theorem 1.1. *Let X be a non-empty set, $x_0 \in X$, Y a real inner product space, $I : X \to \mathbf{R}$, $\Phi : X \to Y$, with $I(x_0) = 0$, $\Phi(x_0) = 0$, and $\mu > 0$. Assume that*

$$\inf_{x\in X} I(x) < 0 \leq \inf_{x\in X}(I(x) + \mu\|\Phi(x)\|^2) \ .$$

Then, for each filtering cover $\mathcal{N}$ of X, there exists $A \in \mathcal{N}$ such that

$$\sup_{y\in Y}\inf_{x\in A}(I(x) + \mu(2\langle\Phi(x), y\rangle - \|y\|^2)) < \inf_{x\in A}\sup_{y\in\Phi(A)}(I(x) + \mu(2\langle\Phi(x), y\rangle - \|y\|^2)) \ .$$

Proof. The assumptions imply that x_0 is a global minimum of $x \to I(x) + \mu\|\Phi(x)\|^2$. But, at the same time, since $\inf_X I < 0$, x_0 is not a global minimum of I. Hence, (b) of Theorem A does not hold and so (a) holds. △

Theorem 1.2. *Let X be a non-empty symmetric set in a real vector space, Y a real inner product space, $I : X \to \mathbf{R}$ an even function and $\Phi : X \to Y$ an odd function.*

Then, for each $\mu > 0$, at least one of the following assertions holds:
(a_1) *for each filtering cover $\mathcal{N}$ of X, there exists $A \in \mathcal{N}$ such that*

$$\sup_{y\in Y}\inf_{x\in A}(I(x) + \mu(2\langle\Phi(x), y\rangle - \|y\|^2)) < \inf_{x\in A}\sup_{y\in\Phi(A)}(I(x) + \mu(2\langle\Phi(x), y\rangle - \|y\|^2)) \ ;$$

(b_1) *if $u \in X$ is a global minimum of $x \to I(x) + \mu\|\Phi(x)\|^2$, then $\Phi(u) = 0$ and u is a global minimum of I.*

Proof. Assume that (a_1) does not hold. Let $u \in X$ be any global minimum of $x \to I(x) + \mu\|\Phi(x)\|^2$. Then, by Theorem A, we have

$$I(u) \leq I(x) + 2\mu(\langle\Phi(x), \Phi(u)\rangle - \|\Phi(u)\|^2) \tag{1}$$

for all $x \in X$. So, in particular, we have

$$I(u) \leq I(-u) + 2\mu(\langle\Phi(-u), \Phi(u)\rangle - \|\Phi(u)\|^2) = I(u) - 4\mu\|\Phi(u)\|^2$$

which yields

$$\|\Phi(u)\|^2 = 0 \ .$$

So, by (1), we get

$$I(u) \leq I(x)$$

for all $x \in X$. △

The application of Theorem 1.1 we wish to present is provided by the following result jointly with Theorem 1.4:

Theorem 1.3. *Let X be a real inner product space and let τ be a topology on X. Moreover, let $J : X \to \mathbf{R}$ be a functional such that*

$$J(0) = 0 < \sup_X J$$

and

$$\beta^* := \sup_{x\in X\setminus\{0\}} \frac{J(x)}{\|x\|^2} < +\infty \ . \tag{2}$$

Finally, let $\lambda > \frac{1}{\beta^*}$ and let $\mathcal{N}$ be a filtering cover of X such that, for each $A \in \mathcal{N}$ and each $y \in X$, the restriction to A of the functional $x \to \|x\|^2 - \lambda J(x) + \langle x, y\rangle$ is τ-lower semicontinuous and inf-τ-compact.

Then, there exists $\tilde{A} \in \mathcal{N}$ with the following property: for every convex set $C \subseteq X$ whose closure (in the strong topology) contains $\tilde{A}$, there exists $\tilde{y} \in C$ such that the restriction to $\tilde{A}$ of the functional $x \to \|x\|^2 - \lambda J(x) + \langle x, 2(\beta^*\lambda - 1)\tilde{y}\rangle$ has at least two global minima.

Proof. In view of (2), we have

$$\inf_{x\in X}(\|x\|^2 - \lambda J(x)) < 0 \ ,$$

as well as

$$\inf_{x\in X}(\|x\|^2 - \lambda J(x) + (\beta^*\lambda - 1)\|x\|^2) \geq 0 \ .$$

So, we can apply Theorem 1.1 taking $Y = X$,

$$\mu = \beta^*\lambda - 1 \ ,$$

$$I(x) = \|x\|^2 - \lambda J(x)$$

and

$$\Phi(x) = x \ .$$

Therefore, there exists $\tilde{A} \in \mathcal{N}$ such that

$$\sup_{y\in Y}\inf_{x\in\tilde{A}}(\|x\|^2 - \lambda J(x) + (\beta^*\lambda - 1)(2\langle x, y\rangle - \|y\|^2)) < \inf_{x\in\tilde{A}}\sup_{y\in\tilde{A}}(\|x\|^2 - \lambda J(x) + (\beta^*\lambda - 1)(2\langle x, y\rangle - \|y\|^2)) \ . \quad (3)$$

Now, consider the function $f : X \times X \to \mathbf{R}$ defined by

$$f(x, y) = \|x\|^2 - \lambda J(x) + (\beta^*\lambda - 1)(2\langle x, y\rangle - \|y\|^2)$$

for all $(x, y) \in X \times X$. Since $f(x, \cdot)$ is continuous and $\tilde{A} \subseteq \overline{C}$, we have

$$\sup_{y\in\tilde{A}} f(x, y) = \sup_{v\in\overline{\tilde{A}}} f(x, y) \leq \sup_{v\in\overline{C}} f(x, y) = \sup_{y\in C} f(x, y)$$

for all $x \in X$, and hence, taking (3) into account, it follows that

$$\sup_{y\in C}\inf_{x\in\tilde{A}} f(x, y) < \inf_{x\in\tilde{A}}\sup_{y\in\tilde{A}} f(x, y) \leq \inf_{x\in\tilde{A}}\sup_{y\in C} f(x, y) \ . \quad (4)$$

Now, in view of (4), taking into account that $f_{|\tilde{A}\times C}$ is τ-lower semicontinuous and inf-τ-compact in $\tilde{A}$, and continuous and concave in C, we can apply Theorem 3.2 of [8] to $f_{|\tilde{A}\times C}$. Consequently, there exists $\tilde{y} \in C$ such that $f_{|\tilde{A}}(\cdot, \tilde{y})$ has at least two global minima, and the proof is complete. △

From Theorem 1.3, in turn, we obtain the following result (with the convention $\frac{1}{0} = +\infty$):

Theorem 1.4. *Let X be a real Hilbert space and let $J : X \to \mathbf{R}$ be a C^1 functional, with compact derivative, such that*

$$\alpha^* := \max\left\{0, \limsup_{\|x\|\to+\infty}\frac{J(x)}{\|x\|^2}\right\} < \beta^* := \sup_{x\in X\setminus\{0\}}\frac{J(x)}{\|x\|^2} < +\infty \ .$$

Then, for every $\lambda \in \left]\frac{1}{2\beta^*}, \frac{1}{2\alpha^*}\right[$ and for every convex set $C \subseteq X$ dense in X, there exists $\tilde{y} \in C$ such that the equation

$$x = \lambda J'(x) + \tilde{y}$$

has at least three solutions, two of which are global minima of the functional $x \to \frac{1}{2}\|x\|^2 - \lambda J(x) - \langle x, \tilde{y}\rangle$.

Proof. Fix $\lambda \in \left]\frac{1}{2\beta^*}, \frac{1}{2\alpha^*}\right[$ and a convex set $C \subseteq X$ dense in X. For each $y \in X$, we have

$$\liminf_{\|x\|\to+\infty}\left(1 - 2\lambda\frac{J(x)}{\|x\|^2} - \frac{\langle x, y\rangle}{\|x\|^2}\right) = 1 - 2\lambda\limsup_{\|x\|\to+\infty}\frac{J(x)}{\|x\|^2} > 0 \ .$$

So, from the identity

$$\|x\|^2 - 2\lambda J(x) - \langle x, y\rangle = \|x\|^2 \left(1 - 2\lambda \frac{J(x)}{\|x\|^2} - \frac{\langle x, y\rangle}{\|x\|^2}\right)$$

it follows that

$$\lim_{\|x\|\to+\infty} (\|x\|^2 - 2\lambda J(x) - \langle x, y\rangle) = +\infty \ . \tag{5}$$

Since J' is compact, J is sequentially weakly continuous ([12], Corollary 41.9). Then, in view of (5) and of the Eberlein-Smulyan theorem, for each $y \in X$, the functional $x \to \|x\|^2 - 2\lambda J(x) + \langle x, y\rangle$ is inf-weakly compact in X. So, we can apply Theorem 1.3 taking the weak topology as τ and $\mathcal{N} = \{X\}$. Consequently, since the set $\frac{1}{1-2\beta^*\lambda}C$ is convex and dense in X, there exists $\hat{y} \in \frac{1}{1-2\beta^*\lambda}C$ such that the functional $x \to \|x\|^2 - 2\lambda J(x) + \langle x, 2(2\beta^*\lambda - 1)\hat{y}\rangle$ has at least two global minima in X which are two of its critical points. Since the same functional satisfies the Palais-Smale condition ([12], Example 38.25), it has a third critical point in view of Corollary 1 of [1]. Clearly, the conclusion follows taking $\tilde{y} = (1 - 2\beta^*\lambda)\hat{y}$. △

We now give a specific application of Theorem 1.4.

Let $\Omega \subset \mathbf{R}^n$ be a bounded domain with smooth boundary. On the Sobolev space $H_0^1(\Omega)$, we consider the scalar product

$$\langle u, v\rangle = \int_\Omega \nabla u(x)\nabla v(x)dx$$

with the induced norm

$$\|u\| = \left(\int_\Omega |\nabla u(x)|^2 dx\right)^{\frac{1}{2}} \ .$$

We denote by $H^{-1}(\Omega)$ the dual of $H_0^1(\Omega)$.

If $n \geq 2$, we denote by $\mathcal{A}$ the class of all Carathéodory functions $f : \Omega \times \mathbf{R} \to \mathbf{R}$ such that

$$\sup_{(x,\xi)\in\Omega\times\mathbf{R}} \frac{|f(x,\xi)|}{1 + |\xi|^q} < +\infty \ ,$$

where $0 < q < \frac{n+2}{n-2}$ if $n > 2$ and $0 < q < +\infty$ if $n = 2$. While, when $n = 1$, we denote by $\mathcal{A}$ the class of all Carathéodory functions $f : \Omega \times \mathbf{R} \to \mathbf{R}$ such that, for each $r > 0$, the function $x \to \sup_{|\xi|\leq r} |f(x,\xi)|$ belongs to $L^1(\Omega)$.

Given $f \in \mathcal{A}$ and $\varphi \in H^{-1}(\Omega)$, consider the following Dirichlet problem

$$\begin{cases} -\Delta u = f(x,u) + \varphi & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \ . \end{cases} \tag{$P_{f,\varphi}$}$$

Let us recall that a weak solution of $(P_{f,\varphi})$ is any $u \in H_0^1(\Omega)$ such that

$$\int_\Omega \nabla u(x)\nabla v(x)dx - \int_\Omega f(x,u(x))v(x)dx - \varphi(v) = 0$$

for all $v \in H_0^1(\Omega)$.

Let $\Psi, J_f : H_0^1(\Omega) \to \mathbf{R}$ be the functionals defined by

$$\Psi(u) = \frac{1}{2}\|u\|^2$$

$$J_f(u) = \int_\Omega F(x,u(x))dx \ ,$$

where

$$F(x,\xi) = \int_0^\xi f(x,t)dt \ .$$

Notice that J_f is a C^1 functional whose derivative is given by

$$J'_f(u)(v) = \int_\Omega f(x, u(x))v(x)dx$$

for all $u, vH_0^1(\Omega)$. Consequently, the weak solutions of problem $(P_{f,\varphi})$ are exactly the critical points in $H_0^1(\Omega)$ of the functional $\Psi - J_f - \varphi$. Moreover, J'_f is compact.

Furthermore, if $f \in \mathcal{A}$ and the set

$$\left\{ x \in \Omega : \sup_{\xi \in \mathbf{R}} F(x, \xi) > 0 \right\}$$

has a positive measure, we have $\sup_{H_0^1(\Omega)} J_f > 0$ (see the proof of Theorem 2 of [9]).

We now state the following

Theorem 1.5. *Let $f \in \mathcal{A}$ be such that the set*

$$\left\{ x \in \Omega : \sup_{\xi \in \mathbf{R}} F(x, \xi) > 0 \right\}$$

has a positive measure and

$$\limsup_{|\xi| \to +\infty} \frac{\sup_{x \in \Omega} F(x, \xi)}{\xi^2} \leq 0 \ . \tag{6}$$

Then, for every $\lambda > 0$ large enough and for every convex set $C \subset H^{-1}(\Omega)$ dense in $H^{-1}(\Omega)$, there exists $\varphi \in C$ such that the problem

$$\begin{cases} -\Delta u = \lambda f(x, u) + \varphi & \text{in } \Omega \\ \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

has at least three weak solutions, two of which are global minima in $H_0^1(\Omega)$ of the functional $\Psi - \lambda J_f - \varphi$.

Proof. Condition (6) clearly implies that

$$\limsup_{\|u\| \to +\infty} \frac{J_f(u)}{\|u\|^2} \leq 0 \ .$$

Then, in view of the above-recalled preliminaries, we can apply Theorem 1.4 taking $X = H_0^1(\Omega)$, $J = J_f$, and the conclusion directly follows. △

The last result is an application of Theorem 1.2.

Let $(T, \mathcal{F}, \mu)$ be a measure space, E a real Banach space and $p \geq 1$.

As usual, $L^p(T, E)$ denotes the space of all (equivalence classes of) strongly μ-measurable functions $u : T \to E$ such that $\int_T \| u(t) \|^p \, d\mu < +\infty$, equipped with the norm

$$\| u \|_{L^p(T,E)} = \left(\int_T \| u(t) \|^p \, d\mu \right)^{\frac{1}{p}} \ .$$

A set $D \subseteq L^p(T, E)$ is said to be decomposable if, for every $u, v \in D$ and every $A \in \mathcal{F}$, the function

$$t \to \chi_A(t)u(t) + (1 - \chi_A(t))v(t)$$

belongs to D, where χ_A denotes the characteristic function of A.

Theorem 1.6. *Let $(T,\mathcal{F},\mu)$ be a non-atomic measure space, with $0<\mu(T)<+\infty$, and E a real Banach space. Let $f:E\to\mathbf{R}$ be a lower semicontinuous and even function which has no global minima. Let $g:T\times E\to\mathbf{R}$ be a Carathéodory function such that $g(x,\cdot)$ is odd for all $t\in T$. Moreover, assume that*

$$\max\left\{\sup_{x\in E}\frac{|f(x)|}{1+\|x\|^p},\sup_{(t,x)\in T\times E}\frac{|g(t,x)|}{1+\|x\|^p}\right\}<+\infty$$

for some $p\geq 1$. For each $u\in L^p(T,E)$, set

$$J(u)=\int_T f(u(t))d\mu+\left(\int_T g(t,u(t))d\mu\right)^2 .$$

Then, the restriction of the functional J to any symmetric and decomposable subset of $L^p(T,E)$ containing the constant functions has no global minima.

Proof. Let $X\subseteq L^p(T,E)$ be a symmetric and decomposable set containing the constant functions. Arguing by contradiction, assume that $\hat{u}\in X$ is a global minimum of $J_{|X}$. For each $h\in L^p(T)$, set

$$X_h=\{u\in X:\|u(t)\|\leq h(t)\text{ a.e. in T}\} .$$

Clearly, X_h is decomposable and the family $\{X_h\}_{h\in L^p(T)}$ is a filtering covering of X. Let $I,\Phi:L^p(T,E)\to\mathbf{R}$ be the functionals defined by

$$I(u)=\int_T f(u(t))d\mu$$

and

$$\Phi(u)=\int_T g(t,u(t))d\mu$$

for all $u\in L^p(T,E)$. Of course, I is lower semicontinuous and even, while Φ is continuous and odd. Fix $h\in L^p(T)$. Notice that X_h is connected ([11]). Therefore, $\Phi(X_h)$ is an interval, and so

$$\Lambda_h:=\overline{\Phi(X_h)}$$

is a compact interval. Now, we can apply Theorem 1.D of [6]. Thanks to it, we then have

$$\sup_{\lambda\in\Lambda_h}\inf_{u\in X_h}(I(u)+2\lambda\Phi(u)-\lambda^2)=\inf_{u\in X_h}\sup_{\lambda\in\Lambda_h}(I(u)+2\lambda\Phi(u)-\lambda^2) .$$

Hence, (with $\mu=1$) assertion (a_1) of Theorem 1.2 does not hold, and so assertion (b_1) must hold. Therefore, $\hat{u}$ is a global minimum of $I_{|X}$ and $\Phi(\hat{u})=0$. Since X contains the constants, we have

$$I(\hat{u})\leq\mu(T)\inf_E f .$$

This clearly implies that

$$\int_T(f(\hat{u}(t))-\inf_E f)d\mu=0 ,$$

and so $\hat{u}(t)$ would be a global minimum of f for μ-almost $t\in T$, against the assumptions. △

Acknowledgement. The author has been supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

REFERENCES

[1] P. Pucci and J. Serrin, A mountain pass theorem, *J. Differential Equations*, 60:142-149, 1985.

[2] B. Ricceri, Some topological mini-max theorems via an alternative principle for multifunctions, *Arch. Math.* (Basel), 60:367-377, 1993.

[3] B. Ricceri, On a topological minimax theorem and its applications, in "Minimax theory and applications", B. Ricceri and S. Simons eds., Kluwer Academic Publishers, 191-216, 1998.

[4] B. Ricceri, A further improvement of a minimax theorem of Borenshtein and Shul'man, *J. Nonlinear Convex Anal.*, 2:279-283, 2001.

[5] B. Ricceri, Minimax theorems for limits of parametrized functions having at most one local minimum lying in a certain set, *Topology Appl.*, 153:3308-3312, 2006.

[6] B. Ricceri, Recent advances in minimax theory and applications, in "Pareto Optimality, Game Theory and Equilibria", A. Chinchuluun, P.M. Pardalos, A. Migdalas, L. Pitsoulis eds., Springer, 23-52, 2008.

[7] B. Ricceri, *Nonlinear eigenvalue problems*, in "Handbook of Nonconvex Analysis and Applications" D. Y. Gao and D. Motreanu eds., International Press, 543-595, 2010.

[8] B. Ricceri, A strict minimax inequality criterion and some of its consequences, *Positivity*, 16:455-470, 2012.

[9] B. Ricceri, Energy functionals of Kirchhoff-type problems having multiple global minima, *Nonlinear Anal.*, 115:130-136, 2015.

[10] B. Ricceri, A minimax theorem in infinite-dimensional topological vector spaces, preprint.

[11] J. Saint Raymond, Connexité des sous-niveaux des fonctionnelles intégrales, *Rend. Circ. Mat. Palermo*, 44:162-168, 1995.

[12] E. Zeidler, Nonlinear functional analysis and its applications, vol. III, Springer-Verlag, 1985.